

# Distribution of X-ray Emission from Jet Knots of 3C273

*E.Yu. Bannikova<sup>1,2</sup>, M.S. Mykhailova<sup>2</sup>*

<sup>1</sup>Radio Astronomy Institute NAS of Ukraine

<sup>2</sup>Karazin Kharkov National University

bannikova@astron.kharkov.ua, aniramtiger@rambler.ru

The jet of the quasar 3C273 is observed at different spectral bands. This jet has a knot structure. Jet radiation in radio and optical bands are connected with synchrotron mechanism, while the emission mechanism producing the X-rays is controversial. We suppose that the X-rays observed for two knots nearest to the quasar can originate from the inverse Compton scattering of external source radiation on relativistic electrons. But in the jet region with constant low X-ray intensity the inverse Compton scattering on cosmic microwave background photons (IC/CMB) is essential because the energy density of external source decreases. Upon this scenario the constraints on the angle between the jet axis and line sight values have been obtained. Also, some physical parameters for two nearest knots of the jet of 3C273 have been estimated.

## Introduction

The jet of 3C273 was observed with high resolution by MERLIN [3], by the Hubble space telescope [1] and Chandra [6] at the radio, optical and X-ray bands correspondingly. Through this work  $H_0 = 75$  km/s/Mpc, so  $1''$  corresponds to 2.37 kpc at the distance of 3C273 [1].


## Jet morphology

In the radio band the jet indicates a 'cocoon structure'. The intensity in the jet increases in the direction away from 3C273. In the optical band  $10''$ -jet is most narrow. Within this jet knots that are location shocks occur where electrons are accelerated at relativistic speeds. Shocks can arise by Kelvin-Helmholtz instability at the boundary between jet matter and ambient. These knots have almost identical brightness and size  $0.1'' - 0.5''$ . The first knot is detected at the distance  $13''$  from the core. To indicate the knots we used notation introduced into practice by Bahcall et al. [1]. The knots A3, In1, In2, Ex1 might not belong to the jet because they are not detected in the radio and X-ray jet images [1]. In the X-ray band the jet has length of  $8''$ . Only two optical knots can be found in it. The remaining jet region has low constant X-ray intensity (see Fig.1).

## Suggested idea

The possibilities for the origin of the X-rays might be [6, 4]: 1) synchrotron emission from a second, more energetic population of electron; 2) inverse Compton scattering (IC) of photons external to the jet; 3) IC of radio-optical synchrotron photons. These theories cannot explain why the ratio of X-ray to radio observed intensities decreases with distance from 3C273. But this fact can be important for study of the jet conditions. And so, in our work it is mostly crucial.

The most plausible mechanism of X-ray emission is IC. The observed intensity distribution was taken from work [5]. It shows that two X-ray intensity peaks correspond to the nearest to the quasar optical knots, A1 and B1. Then this intensity decreases to a lower level but it is still detectable until it reaches the terminating optical knot. While in the optical band the knots have almost identical intensities. Thus, the idea of explaining the X-ray intensity distribution observed along the jet of 3C273 is the following [2]. In different jet regions scattering photons have different origin. For the nearest to the quasar jet region there is



The optical profile and X-ray histogram were scaled to  $0.325 \mu\text{Jy}$  at vertical value of 500. The raw optical profile was smoothed with a Gaussian with  $\sigma=0.25''$  and was scaled to  $0.25 \mu\text{Jy}$  at a vertical value of 500.

Figure 1: Distribution of the X-ray and optical emission along the axis of the jet of 3C273 [5].

a heterogeneous radiation field. While for the remaining jet the homogeneous radiation field predominates. In this case, X-ray emission of the two nearest knots originates from IC of quasar photons. For the remaining knots X-rays arise from the same scattering of cosmic microwave background photons. Thus, the basic term was formulated (1):

$$\begin{cases} W_{ES} > W_{CMB}, & r_{arc} < 16'', \\ W_{ES} < W_{CMB}, & r_{arc} > 16'', \end{cases} \quad (1)$$

where  $W_{ES}$  is the energy density of quasar emission,  $W_{CMB}$  is the energy density of cosmic microwave background,  $r_{arc}$  is the project distance from the quasar in arcseconds,  $16''$  was taken because after this distance from 3C273 the X-ray emission have almost constant value (see Fig.1).

Using the energy densities, the integral values can be considered. The observed intensity decline from the knot A1 to knot B1 might be due to the fact that the ratio of electron concentration in these knots is  $N_A/N_B = 2$ . Electron concentration in remaining knots does not change. Taking into account that the X-ray intensity can be defined, as:

$$I_X \propto N \frac{L}{4\pi R^2}, \quad (2)$$

where  $R$  is the physical distance from the quasar, and knowing the energy density of the cosmic microwave background  $W_{CMB}$  and the ratios of observed intensities at jet knots,  $I_A/I_B \approx 2$ ,  $I_A/I_C \approx 8$ , hence, the energy density of quasar emission at the knot A1 was found:

$$W_{ES}^A = 8 \cdot W_{CMB} \cdot \frac{1}{2}.$$

The distance  $R$  from the knot A1 to 3C273 being known, the quasar luminosity  $L$  has been obtained  $L = 4\pi c R^2 W_{ES}^A$ . It is consistent with the observed one. Hence, this assumption is believable.

## Determination of the jet parameters

Jets enclose the thermal plasma that provides the main contribution to the jet mass. But due to its low concentration it cannot be detected directly. The collisions between particles of the thermal plasma can provoke absorption of the emission. The absorption coefficient is

$$\mu \propto \frac{\omega_p^2 \nu}{\omega c}, \quad (3)$$

where  $\omega_p = \sqrt{\frac{4\pi n e^2}{m}}$  is the plasma frequency,  $n$  and  $e$  are the concentration and the charge of electrons,  $\nu$  is the effective collision frequency.

In this case the energy density of quasar emission is:

$$W_{ES} = \frac{L}{4\pi c r^2} \sin^2 \theta \exp[-\mu \frac{(r - r_0)}{\sin \theta}], \quad (4)$$

where  $r_0$  is the project distance from the quasar to jet begining,  $r$  is the project distance from the quasar,  $\theta$  is the angle between jet axis and line of sight.

If the absorption of the quasar emission in the jet does not exist, then the basic term (1) is true for  $\theta = 10^\circ$ . Taking into account that the jet electrons have a Lorenz factor of  $10^3 \div 10^4$ , only the photons having the frequency  $\omega = (10^9 \div 10^{13})$  Hz before scattering can fall into the Chandra frequency range  $\omega_{Ch} = (10^{17} \div 10^{19})$  Hz. The estimates of the lowest temperature and biggest concentration of the thermal electrons have been made when the absorption coefficient tends to zero. We have obtained values for  $n \sim 0.1 \text{ cm}^{-3}$  and  $T > 10^4 \text{ K}$ .

If  $\theta > 10^\circ$ , then the basic term (1) is true when there is absorption of quasar emission along the jet. The absorption coefficient should be of the order of  $10^{-23} \text{ cm}^{-1}$ . Using the (1), (3) the inequality is:

$$\frac{n^2}{\omega^2 T^{3/2} \sin \theta} > 4 \cdot 10^{-23} \frac{c^2}{\text{cm}^6 K^{3/2}}.$$

We found that either  $n > 10 \text{ cm}^{-3}$  for  $T = (10^3 \div 10^4) \text{ K}$  or  $n \sim 1 \text{ cm}^{-3}$  for  $T \sim 10^2 \text{ K}$ . Using (2) and the expressions for energy densities of the quasar emission at the knots A1, B1 and C1

$$\begin{aligned} W_{ES}^A &= \frac{L}{4\pi c a^2} \sin^2 \theta, \\ W_{ES}^B &= \frac{L}{4\pi c (a + b)^2} \sin^2 \theta \exp[-\frac{\mu b}{\sin \theta}], \\ W_{ES}^C &= \frac{L}{4\pi c (a + 2b)^2} \sin^2 \theta \exp[-\frac{2\mu b}{\sin \theta}], \end{aligned}$$

(where  $a$  is the distance from 3C273 to knot A1,  $b$  is the distance from the knot A1 to knot B1 and the distance between knots B1 and C1 (see Fig.1)), the following inequalities were obtained:

$$\begin{aligned} \sin^2 \theta &> W_{CMB} \frac{4\pi c a^2}{L} \left( \frac{I_A}{I_B} \frac{N_A}{N_B} \right), \\ \sin^2 \theta &< W_{CMB} \frac{4\pi c a^2}{L} \left( \frac{I_B}{I_A} \frac{N_A}{N_B} \right). \end{aligned}$$

They mean that the energy density of quasar emission appears to predominate at the knot B1, while the energy density of cosmic microwave background appears to predominate at the knot C1. From these inequalities the constraint on  $\theta$  value was found:  $18^\circ < \theta < 37^\circ$ .

## Conclusions

We have investigated the images of the jet of 3C273 in the different spectral bands to determine mechanism of X-ray emission from this jet. Having supposed, that the X-ray emission from the nearest to the quasar knots A1 and B1 are produced by inverse Compton scattering (IC) of quasar photons. The X-rays from the jet region with constant X-ray intensity are due to the same scattering of cosmic microwave background photons. Assuming this the concentration and temperature of the thermal electrons in the jet of 3C273 were estimated. Obtained values are consistent with theoretical ones for such objects. Hence, in the such framework we can estimate not only the angle between the jet axis and the line of sight, but also the concentration and the temperature of thermal electrons in the jet of 3C273. We can perform this by comparing jet images in the X-ray, optical and radio bands.

## Acknowledgements

We are thankful to Victor M. Kontorovich for discussions of this work.

## References

- [1] Bahcall J.N., Kirhakos S., Schneider D.P. ApJ, V. 452, pp. L91-L93 (1995)
- [2] Bannikova E.Yu., Kontorovich V.M. Space Science and Technology, V. 5/6, pp. 153-157 (2003)
- [3] Conway R.G., Garrington S.T., Perley R.A. et al. A&A, V.267, pp. 347-362 (1992)
- [4] Harris D.E., Krawczynski H. ApJ, V.565, pp.244-255 (2002)
- [5] Marshall H.L., Harris D.E., Grimes J.P. et al. ApJ, V. 549, pp. 167-171 (2001)
- [6] Sambruna R.M., Urry C.M., Tavecchio F. et al. ApJ, V.549, pp. 161-165 (2001)